BACK HEALTH

Evaluation and Treatment of Sacroiliac Joint Dysfunction in the Primary Care Setting: A Practical Review

ABSTRACT

Mechanical dysfunction of the sacroiliac joint (SIJ) is an often overlooked, but a common cause of low back pain in the North American adult population. The diagnosis is primarily clinical and requires the exclusion of other potential etiologies of low back pain (LBP). A number of non-surgical treatment options are available for patients with this pathological entity. In cases of persistent, severe SIJ pain refractory to non-operative measures, SIJ fusion may be considered as a surgical intervention.

KEYWORDS: sacroiliitis, sacroiliac dysfunction, sacroiliac joint (SIJ), low back pain (LBP), gluteal pain, SI joint fusion, percutaneous SI joint fixation



Introduction

Although the etiologies are diverse and heterogeneous, low back pain (LBP) remains a leading source of morbidity in North America and a major financial burden on the healthcare system.¹ A common cause of LBP is the sacroiliac joint (SIJ), increasingly recognized as a major source of LBP and disability.² SIJ pain can be severely debilitating, leading to substantial functional impairment and diminished quality of life. Studies show that patients with SIJ dysfunction have a quality of life equivalent to those with advanced hip and knee arthritis and worse than those with chronic obstructive pulmonary disease or mild heart failure.²



Sydney Rucker, BS, Medical University of South Carolina College of Medicine, Charleston, SC



Adrienne Kelly, MD, FRCSC, Orthopaedic Surgeon, Sault Area Hospital, Assistant Professor, Northern Ontario School of Medicine, Sault Ste Marie, ON



David W. Polly, MD, Department of Orthopaedic Surgery, University of Minnesota, Minneapolis, MN, USA



Robert J. Ferdon, MS, Department of Orthopaedics and Physical Medicine, Medical University of South Carolina, Charleston, SC, University of South Carolina School of Medicine, Columbia, SC



Robert A. Ravinsky, MDCM, MPH, FRCSC, Department of Orthopaedics and Physical Medicine, Medical University of South Carolina, Charleston, SC

Pre-test Quiz

The notion that the SIJ is a common pain generator is not new but the joint's relevance as a target of both conservative and surgical treatments has waxed and waned over the last several decades. Nevertheless, a growing body of literature has resulted in improved knowledge of the anatomy, physiology, mechanics and pathology of this previously nebulous part of the musculoskeletal system. Improved understanding of SIJ anatomy and biomechanics has allowed for the development of evidence-based treatment pathways and effective treatment modalities for this condition.

In this review article we aim to provide clear guidance for appropriate identification, clinical and radiographic evaluation of mechanical SIJ dysfunction and provide a framework for treatment of this pathological entity in the primary care setting.

Epidemiology

SIJ dysfunction is a known contributor to LBP, accounting for

Table 1: Risk Factors for Sacroiliac JointDysfunction^{8,9}

Idiopathic

Multigravida w/prior history of vaginal birth

High-energy pelvic trauma

Prior history of iliac crest bone grafting

Prior history of lumbosacral fusion (multilevel)

approximately 15-30% of LBP cases in the outpatient setting.³⁻⁷ Several primary risk factors for mechanical SIJ dysfunction have been identified: 1) multigravida females with a prior history of vaginal birth (likely due in part to the increased levels of estrogen or relaxin during the third trimester of pregnancy resulting in increased ligamentous laxity and trauma to the pelvic floor and SIJ ligaments during childbirth.⁸⁻⁹), 2) a prior history of high-energy pelvic trauma or iliac crest bone grafting, because of the extent of the associated soft tissue injury which can result in pathological motion at this joint.¹⁰), 3) a prior history of lumbosacral fusion, particularly multilevel lumbar fusion is a known risk factor for development of secondary mechanical SIJ dysfunction; the causal mechanism is analogous to that of adjacent segment disease, as several previously mobile segments in the lumbar spine and lumbosacral junction are immobilized, stress transferred to the SIJ substantially increases and results in subsequent degeneration. SIJ pain as a secondary source of discomfort following lumbar fusion is exceedingly common, with studies indicating between 34-43% of patients develop SIJ-related pain postoperatively.¹¹⁻¹⁴ Apart from these risk factors, many cases are idiopathic (see Table 1).

SIJ Anatomy and Biomechanics

The sacroiliac joint, the largest axial joint in the human body, is a complex syndesmosis, symphysis and synovial joint connecting the sacrum to the ilium. The joint plays a critical role in transferring spinal loads to the lower extremities, allowing limited but important motion, typically 1.1-2.2 degrees in flexion-extension, 0.5-8.0 degrees in lateral bending and 0.8-4.0 degrees in axial rotation, with minimal translation.¹⁵⁻¹⁹ While movement in these planes supports daily activity, slight variations in movement patterns can significantly impair function.²⁰

SIJ innervation is variable and involves multiple nerves, including the sacral plexus, superior gluteal nerve, dorsal rami of L4-S4, anterior rami of L2-S2 and the obturator nerve. Histologic studies have identified nociceptors and mechanoreceptors, predominantly within the anterior capsule and cartilage, that are mainly supplied by the dorsal rami of S1-S3. Activation of these nociceptors can cause SIJ pain; thus, they are targeted as a therapeutic focus in interventional pain procedures such as radiofrequency neurotomy and injection therapies aimed at denervating or modulating afferent input to the SIJ.²¹⁻²³

Finite element studies suggest that normal SIJ loads can range from 79-140 Newtons during physiological activities.²⁰ The bone

density of the sacrum plays a critical role in force distribution, being highest near the sacral body and lowest in the sacral ala. Variability in trabecular architecture and density influences load transmission and may be implicated in implant loosening and failure of SIJ fusion procedures.²⁴⁻²⁶ Joint stability is mediated by both form and force closure mechanisms. Form closure relies on the inherent anatomical interlocking of the sacral and iliac surfaces, providing passive stability. Force closure, in contrast, is dynamic and depends on the compression forces generated by surrounding muscles, ligaments and fascial structures.²⁷

Anatomically, the SIJ is unique; it is a synovial joint at the anterior third and a fibrous joint in the posterior two-thirds. The considerable stability is attributed in part to the robust anterior and posterior ligamentous complexes. Extensive ligamentous structures and muscles including the erector spinae, lumbar multifidus, gluteal muscles and muscles of the abdominal wall stabilize the remaining portion.²⁸ While these muscles do not directly produce movement across the SIJ, they aid in pelvic stabilization and load transmission. Decreased muscle thickness in the paraspinal musculature has been associated with SIJ-related pain.²⁹ When stabilizing mechanisms are compromised through trauma, degeneration,

hypermobility or other pathologic mechanisms, joint motion and pain may ensue. For example, SIJ dysfunction following lumbar fusion is one frequently cited mechanism of altered stability; the increased stress across the SIJ contributes to subsequent instability and pain.²⁰

Pathophysiology

Mechanical SIJ dysfunction most commonly arises from repetitive trauma that disrupts normal load transfer across the pelvis, leading to pathological motion and degenerative arthropathy.

Pregnancy, pelvic trauma and prior lumbosacral fusion are all examples of this phenomenon. These mechanical insults initiate a cascade of inflammation, nociceptive activation and pain that may begin insidiously or acutely. Key pain generators include increased ligamentous or capsular tension, extraneous compression or shear forces, myofascial or kinetic chain imbalances, joint hypo- or hypermobility or other forces that disrupt normal load transfer across the pelvis.

Non-mechanical causes of SIJ pathology include inflammatory spondyloarthropathies, crystalline arthridites (such as gout), infection, metabolic bone disease and rarely, neoplasms. Infections are most often caused by *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Cryptococcus* and *Mycobacterium* tuberculosis, particularly in patients with immunosuppression, endocarditis, IV drug use or other predisposing conditions. Among neoplastic causes, metastatic disease is most common, with pelvic involvement occurring in nearly 40% of skeletal metastases, second only to the spine.³⁰

Clinical Evaluation

SIJ dysfunction typically presents as buttock pain or LBP, with referral patterns often mimicking other spinal, pelvic or hip pathologies.³¹ Differential diagnoses include piriformis syndrome, lumbar spinal stenosis, lumbar disc herniation, degenerative disc disease, hip osteoarthritis, gluteal tendinopathies, labral tears, hamstring origin avulsion injuries and sciatic, pelvic, or pudendal nerve entrapment via greater trochanteric, ischiofemoral or posterior femoroacetabular impingement or other aberrant anatomical stressors.32 Differentiating mechanical SIJ dysfunction from sacroiliitis, an inflammatory condition, is essential for selecting an appropriate treatment plan.

Accurate diagnosis of SIJ pathology relies heavily on clinical evaluation. Patients typically report Fortin's sign, defined as pain consistently localized to an area within 1cm inferomedial to the posterior superior iliac spine (PSIS) that may radiate into the buttocks, groin, posterior thigh or even past the knee and into the foot.³³⁻³⁴ This reproducible pattern is a key diagnostic feature that strongly suggests SIJ dysfunction and serves as a clinical pearl in LBP evaluation. Pain above the PSIS may suggest alternative etiologies such as facetmediated pain, superior cluneal nerve entrapment or quadratus lumborum myofascial dysfunction. Pain below the PSIS more commonly reflects piriformis syndrome or middle cluneal nerve entrapment.³² Symptoms often worsen with prolonged standing, sitting or transitional movements. Some patients report relief with a tightfitting belt, likely due to SIJ stabilization (see Table 2).

For pain over the PSIS or other physical presentations suggesting SIJ dysfunction, physical examination should include SIJ-specific provocation tests. Level 1 evidence supports a Clinical Diagnostic Rule for SIJ pain, using 3 or more positive results out of five standardized maneuvers, with dominant examination findings being pain in PSIS area. Provocation tests include the distraction (gapping) test, SI compression test, Patrick's (FABER) test, thigh thrust or posterior shear (POSH), sacral thrust and Gaenslen's (Pelvic torsion) test.³⁵⁻³⁶ These diagnostic criteria are good or better than rules for other lumbar spine conditions, making SIJ diagnosis among the best evidence-based diagnostic rules in

Table 2: Differential Diagnoses³²

Piriformis syndrome

Sciatic nerve irritation/entrapment radiating pain

Ischiofemoral impingement and/or semimembranosus origin avulsion

Conjoined hamstring tendon origin avulsion and/or sacrotuberous ligament insertion avulsion

Pudendal nerve irritation/entrapment

Obturator internus tears or source of pundendal irritation

Adductor magnus origin tendinoisis/avulsion

Greater trochanteric pelvic/sciatic impingement and/or greater trochanteric bursitis

Gluteus maximus origin tendonitis

Sacroiliac joint pain/dysfunction

Gluteus maximus claudication

Posterior femoroacetabular impingement and/or posterior labral tear

Hip-Spine syndrome

LBP classification.³⁵ Meeting the criteria of 3 or more positive exam findings has an 85% positive-predictive value of positive response to diagnostic injection.

Radiographic Evaluation

While imaging plays a role in ruling out alternative diagnoses, no imaging modality has demonstrated reliable diagnostic utility for mechanical SIJ dysfunction, though there is some utility of MRI in the setting of inflammatory sacroiliitis.³⁷ Studies have shown that imaging provides minimal benefit in the management of LBP in the absence of red flag symptoms preceding a trial of non-operative management.³⁸ If symptoms do not improve after 6-8 weeks of non-surgical treatment, lumbar radiographs are indicated. Patients presenting with red flag symptoms, however, warrant immediate imaging.

Plain radiographs

Upright anteroposterior (AP) and lateral lumbar spine radiographs are the initial imaging modality of choice for LBP due to their accessibility and economic value. PA and oblique views of the SIJs themselves can be done but rarely influence therapeutic decision-making.

Plain radiographs often reveal degenerative changes of the SIJ, but such findings are also prevalent among asymptomatic adults over the age of 50.³⁹ This high prevalence of SIJ degeneration in pain-

free individuals complicates efforts to attribute LBP or pelvic pain to the SIJ in symptomatic patients. Evidence suggests that SIJ degeneration is a normal part of aging, with many patients exhibiting significant radiographic findings without reporting associated pain. This incongruity between imaging findings and clinical symptoms has contributed to the challenge of diagnosis and treatment of SIJ related pain.⁴⁰

Computed Tomography (CT)

CT is highly effective in detecting structural changes of the SIJ and subchondral demineralization and erosions of the iliac surface early in an inflammatory disease course. In later stages, CT can reveal ankylosis of the SIJ. Compared to MRI, CT is often less expensive and allows for more rapid acquisition of images, but is limited in its' ability to detect bone marrow edema and very early cartilage abnormalities.⁴¹ CT imaging, though useful in some settings for identifying sacroiliitis or fractures, demonstrates limited sensitivity (57%) and specificity (69%) for mechanical dysfunction.42

Magnetic Resonance Imaging (MRI)

MRI remains the gold standard and most sensitive imaging modality for detecting inflammatory SI pathology manifesting as bone marrow edema, particularly in spondyloarthropathies, but has limited utility in purely mechanical cases.⁴³⁻⁴⁴ MRI of the lumbar spine allows for visualization of degenerative changes within the intervertebral discs and facet joints and identification of spinal stenosis involving the central canal and/or neuroforamina.⁴⁵⁻⁴⁶

Nuclear Medicine

Nuclear medicine imaging techniques, such as bone scintigraphy and gallium scans, may be helpful in specific clinical scenarios, such as detecting malignancies, infections, and certain bone disorders like Paget's disease and fibrous dysplasia. Bone scintigraphy with single photon emission computed tomography (SPECT/CT) is a more recent imaging modality that has been shown to improve anatomical localization of active lesions through radiotracer uptake in patients with non-specific musculoskeletal pain. These modalities, however, generally lack sufficient sensitivity for routine use in the context of SIJ dysfunction.⁴⁷

Diagnostic Injections

Diagnostic intra-articular SIJ injections performed under fluoroscopy or CT guidance are considered the gold standard for confirming SIJ-mediated pain (see Figure 1). Contrast-enhanced imaging ensures accurate placement, addressing the high rates of inaccuracy with blind injections. Rosenberg et al. found that only 22% of blind SIJ injections were within the joint space.⁴⁸⁻⁵⁰



Figure 1: Demonstrates fluoroscopic intra-articular SIJ injection, done in the procedure suite by physiatrist. Image 1A is seen on the left and demonstrates placement of the spinal needle in the synovial portion of the SIJ. Image 1B demonstrates extravasation of radiopaque dye within the SIJ, confirming accurate placement of the needle.

Non-surgical Treatment

Non-surgical treatment remains the primary management approach for SIJ dysfunction. Several conservative treatment modalities have been extrapolated from the LBP literature, given the overlap in symptomatology and biomechanical principles. Pharmacologic therapies including acetaminophen, non-steroidal anti-inflammatories (NSAIDs), muscle relaxants, opioids and topical analgesics may provide symptomatic relief. Bracing and physical therapy may provide reasonable adjuncts. Further treatment options for non-refractory pain include image-guided intra-articular corticosteroid injections and radiofrequency ablation (RFA) of lateral branch nerves.

Pharmaceutical Treatment

The American Geriatric Society guidelines recommend acetaminophen as a first-line treatment for LBP due to its favorable safety profile, but a systematic review concluded acetaminophen was ineffective treating acute LBP; van der Gaag et al. found NSAIDs to be slightly effective in reducing the intensity of acute LBP.⁵¹⁻⁵² Evidence regarding the effectiveness of these over-the-counter analgesics in managing chronic LBP remains conflicting; high-quality evidence from three studies with a total of 1,825 participants suggested that acetaminophen was no more effective than placebo.53

Literature on the utility of muscle relaxants is controversial. A systematic review of 30 trials found benzodiazepines, non-benzodiazepines and anti-spasticity muscle relaxants (in combination with benzodiazepines and nonbenzodiazepines) were effective in the treatment of non-specific LBP. While a systematic review of 49 trials reported no clinically meaningful pain relief.⁵⁴⁻⁵⁵ This uncertainty, coupled with the risk of adverse events calls for caution in the use of muscle relaxants for LBP management.

For patients with chronic LBP, opioids like tramadol produce modest, short-term pain relief that may or may not be clinically meaningful.⁵⁶⁻⁵⁸ Due to the potential for drug dependence and negative side effects, we recommend against the use of opioid medication for SIJrelated pain (see Table 3).

Topicals

Topical analgesics such as lidocaine, diclofenac and capsaicin have shown some promise in reducing non-specific LBP alone or in combination with systemic analgesics.⁵⁹⁻⁶¹ While evidence for their efficacy when applied to SIJ pain is limited, the low-risk side-effect profile makes them a reasonable alternative to systemic pharmacologic options.⁶²

Bracing

The use of pelvic belts in combination with muscle strengthening may increase pelvic stability by reducing the sagittal rotation in the SIJs and alleviating tension in the SIJ ligaments.⁶³⁻⁶⁴

Weight Reduction

The specific data considering the relationship between obesity/ elevated body mass index and mechanical SIJ pain is sparse; however, given that the SIJ is a load-bearing joint, an association between obesity and LBP seems

Table 3: Conservative Treatment Options^{8,63}

Medications

- Acetaminophen/Paracetamol
- NSAIDs
- Muscle relaxants
- Tramadol
- Oral steroids
- Atypical anticonvulsants

Topicals

- Lidocaine
- Diclofenac
- Capsaicin

Physical therapy (PT)

Bracing

Pelvic belts

Weight loss

Image guided injections

- CT-guided corticosteroid injection
- Ultrasound guided corticosteroid injection

Radiofrequency ablation (RFA)

• Lateral branch nerves of S1, S2, S3

plausible. Evidence suggests a link between obesity and LBP, likely due to increased mechanical stress on the spine and systemic inflammation. The same is likely true with the SIJ, but there is limited research confirming that weight loss significantly reduces LBP symptoms.⁶⁵ A systematic review of surgical and non-operative weight loss interventions found only lowquality evidence of improvements in pain, disability or quality of life.66 While weight loss may offer some benefit for SIJ pain, current evidence supports moderate weight reduction to be used in conjunction with other non-operative treatment modalities.

Physical Therapy

Structured physical therapy programs that emphasize core strengthening, pelvic stabilization and manual therapy techniques may be beneficial for SIJ pain. During the acute phase, deficits in flexibility and strength should be identified and addressed to minimize the effects of reduced mobility, including muscle atrophy and ligament weakening.8 Given that deconditioning of the paraspinal muscles has been implicated in the pathogenesis of mechanical SIJ pain, physical therapy directed at improved core strength and muscular control is a logical initial option for patients with presumptive mechanical SIJ dysfunction.

Image Guided Injections

In the management of SIJ pain, image-guided corticosteroid injections may be considered when oral medications and physical therapy fail to provide adequate pain relief.⁶⁷ Image guidance may be achieved via several different modalities, such as fluoroscopy, ultrasound or CT. Image-guided corticosteroid injections serve both diagnostic and therapeutic purposes, although their effects may in some instances be temporary. Nevertheless, injections are often a useful treatment, particularly when the etiology of the pain is inflammatory. One study found that CT-guided corticosteroid injections resulted in a significant reduction in inflammatory back pain and sacroiliitis in patients with spondyloarthropathies.68 Conversely, in cases of hypermobility or mechanical dysfunction, steroid injections may not afford durable relief.

Radiofrequency Ablation (RFA)

For patients who have SIJ pain refractory to non-operative measures, RFA of lateral branch nerves sacral nerves, innervating the SIJ has demonstrated short-term pain relief in some studies. RFA of the lateral branch nerves as a treatment for SIJ pain is particularly helpful in patients in whom nociceptive fibers have been pathologically activated. One meta-analysis found that at 3- and 6-month intervals, 60.1% and 49.9% of patients, respectively, experienced at least 50% pain relief. The diminished outcomes over time were likely attributable to the nerve regeneration and regrowth.⁶⁹

Overall, non-operative management is safe and typically associated with some reduction in pain although durability of relief may be variable. There are limitations. This approach is associated with substantial costs of care and may provide limited long-term benefit;⁷⁰⁻⁷² however, some patients certainly benefit from non-operative care and our recommendation is for a thorough course be attempted in all patients prior to the consideration of surgery.

Surgical Treatment

Surgical intervention may be considered for patients with clinically debilitating mechanical SIJ dysfunction who fail exhaustive non-operative therapy, though access remains limited in some regions.

Historically, external frame fixation was explored, but proved impractical and excessively morbid. This approach is not a currently accepted practice and is of academic interest only.⁷³ More recently, procedures to stabilize and fuse the SIJ have been developed using implants such as triangular titanium rods (TTRs), porous fusion/fixation screws (PFFS) and sacropelvic fixation strategies. The most extensively studied technique is percutaneous SIJF, involving a lateral percutaneous approach with placement of porous triangular titanium implants across the synovial portion of the SIJ (see Figure 2). This method is gaining widespread traction as



Figure 2: Postoperative films in a 64-year-old male that underwent percutaneous SIJF. The figure demonstrates AP (2A), pelvic inlet (2B), pelvic outlet and lateral views of the pelvis and sacrum. This patient, with a prior history of multilevel lumbosacral fusion presented with clinical complaints related to significant, debilitating left-sided buttock pain, as well as several positive provocative physical examination findings suggestive of SIJ dysfunction. He was investigated thoroughly, and other possible pain generators in the differential diagnosis were systematically ruled out. He then underwent 2 fluoroscopically guided intra-articular SIJ injections. The patient had a brief but profound response. He was subsequently electively booked for percutaneous left-sided SIJ fusion with use of porous triangular implants (SI-BONE). After surgery, he experienced very significant relief of left-sided buttock pain. At long-term follow-up he was experiencing minimal pain and had a dramatically improved level of physical function.

an evidence-based treatment, supported by Level I evidence, including two randomized controlled trials, multiple prospective cohort studies and over 100 peer-reviewed publications. Multiple prospective clinical trials have consistently demonstrated significant improvements in pain and disability scores.^{70,74-77} Typically performed in the outpatient setting, this procedure has been associated with reduced opioid consumption, improved returnto-work rates, high patient satisfaction and sustained pain relief at 5 years. Operating times average 45-60 minutes, with minimal blood loss (see Figure 3). The technique has been shown to be cost-effective, with an estimated incremental cost effectiveness ratio (ICER) of \$13,000 USD, comparable to total knee arthroplasty (ICER \$12,000).⁷¹ Complications include surgical site infection, gluteal pain, implant loosening or failure, mispositioning, pseudoarthrosis, nerve injury and persistent pain; how-



Figure 3: SIJF as a component of adult spinal deformity correction with fusion to pelvis. This 57-year-old female had previously undergone T9-L3 instrumented fusion for infection. After the infection had cleared, the patient was experiencing worsening back pain, bilateral buttock pain and difficulty standing upright. She was determined to have a spinal deformity that was amenable to surgical correction. She underwent revision deformity correction, and as part of fusion construct multiple screws were placed into the pelvis, including 2 porous titanium screws, which were inserted from the sacrum into the pelvis, across the SIJ. The inherent design of the screws is meant to allow for SIJF. Furthermore, the use of more than one point of fixation across the SIJ allows for rotational stabilization of this joint.

ever, complication rates are low overall and the safety profile for these procedures is generally favorable, with revision rates decreasing over time.⁷⁷⁻⁷⁸ A recent meta-analysis confirmed that minimally invasive SIJF, particularly the lateral transiliac (LTI) approach, results in significant improvement in pain and disability, with low adverse event rates. In contrast, posterolateral (PLTI) and posterior intra-articular (PI) approaches are less studied and have less predictable results.⁷⁹

Surgical SIJF may be beneficial for patients undergoing spinal deformity correction surgery or long lumbar fusions to the sacrum in pelvis. These conditions include significant deformity corrections, fusions of L2 or higher to sacrum, lumbosacral fusions in patients with a diagnosis of SIJ pain and lumbosacral fusion in patients with increased BMI. In these patients, mechanical stresses imparted to the SIJ substantially increases, resulting in an increased risk of subsequent SIJ dysfunction. Fusing the SIJ may prevent pelvic screw loosening or rod fracture, because there is a high rate of mechanical complications in these patients at long-term follow up.⁸⁰⁻⁸²

Defining ideal surgical candidates remains a challenge. Current criteria include \geq 3 positive SIJ provocative tests, diagnostic pain relief from two image-guided SIJ injections and exclusion of other pain sources. Thresholds for diagnostic response to SIJF vary from 50-90% pain relief; 75% is most commonly used, but some suggest 50% may be more appropriate in patients with multiple pain generators, such as spinal degenerative disease.^{9,21,31,49,50} Polly et al. cautioned against overly stringent thresholds (75% or higher) to determine surgical candidacy, because this may exclude patients who would otherwise benefit from surgical intervention.

From a broader perspective, there may be some patients with multiple sources of pain, in whom

- 🖸 SUMMARY OF KEY POINTS -

- 1. The SI joint is a known contributor to low back pain
- 2. Non-surgical treatment remains the primary management approach for SIJ dysfunction
- 3. Diagnostic intra-articular injections performed under image guidance are considered the gold standard for confirming SIJ-mediated pain
- 4. For patients with clinically debilitating mechanical SIJ dysfunction who have failed an exhaustive course of nonoperative treatment, surgical options may be considered

SIJF may afford significant pain relief; however, these patients may describe a lower percentage of pain relief than the previously described 75% threshold. In a subgroup analysis of 320 subjects from multiple centers who failed non-surgical management, authors found that the degree of acute pain relief from diagnostic SIJ injections did not correlate with improved pain or function at 6 or 12 months post-SIJF.⁸¹ As such, ongoing refinements of clinical cut-off points for surgical eligibility require further investigation as do clinical outcomes before widespread adoption of these surgical techniques are seen.

Future Directions

Future research directions include the ongoing SILVIA randomized controlled trial, evaluating concurrent SIJF during multilevel spine reconstructions. Despite growing evidence supporting isolated percutaneous SIJF, further study is needed to refine patient selection criteria, and determine best practices for patients with complex spinopelvic pathologies. This approach remains an active area of investigation.

Conclusions

SIJ dysfunction is an underdiagnosed yet frequent cause of LBP. Careful clinical evaluation, supported by diagnostic injections, is critical for accurate identification. While non-surgical therapies are appropriate initial strategies, some patients may fail an exhaustive course and ultimately may benefit from SIJF. As our understanding of SIJ pathology continues to expand, the optimization of diagnostic and therapeutic pathways promises to further improve outcomes for treating patients with this SIJ-related pain.



CLINICAL PEARLS

- 1. SIJ dysfunction is a known contributor to low back pain, accounting for approximately 15-30% of LBP cases in the outpatient setting.
- 2. While imaging plays a role in excluding alternative diagnoses, no imaging modality has demonstrated reliable diagnostic utility for mechanical SIJ dysfunction although there is some utility of MRI in the setting of inflammatory sacroiliitis.
- 3. Patients with SIJ pain typically report symptoms consistently localized to an area within 1cm inferomedial to the posterior superior iliac spine (PSIS) that may radiate into the buttocks, groin, posterior thigh or even past the knee and into the foot.
- 4. Diagnosis is by physical examination which should include a variety of SIJ-specific provocation tests. Three or more positive results out of five standardized maneuvers is supported by Level 1 evidence for a Clinical Diagnostic Rule.

сме Post-test Quiz

Members of the College of Family Physicians of Canada may claim MAINPRO-M2 Credits for this unaccredited educational program.

References

- 1. Kirsch EP, Yang S, Li Y, et al. Healthcare resource utilization for chronic low back pain among high-utilizers. Spine J. 2024;24(4):601-616
- 2. Cher DJ, Polly DW, Berven SH. Sacroiliac joint pain: burden of disease. Med Devices (Auckl). 2014;7:73-81. doi:10.2147/MDER.S61821
- 3. Bernard TN Jr, Kirkaldy-Willis WH. Recognizing specific characteristics of nonspecific low back pain. Clin Orthop Relat Res. 1987;(217):266-280.
- 4. Schwarzer AC, Aprill CN, Bogduk N. The sacroiliac joint in chronic low back pain. Spine (Phila Pa 1976). 1995;20(1):31-37. doi:10.1097/00007632-199501000-00007.
- 5. Maigne JY, Aivaliklis A, Pfefer F. Results of sacroiliac joint double block and value of sacroiliac pain provocation tests in 54 patients with low back pain. Spine (Phila Pa 1976). 1996;21(16):1889-1892. doi:10.1097/00007632-199608150-00010.
- 6. Irwin RW, Watson T, Minick R, Ambrosius WT. Age, body mass index, and gender differences in sacroiliac joint pathology. Am J Phys Med Rehabil. 2007;86(1):37-44. doi:10.1097/ PHM.0b013e31802b8723.
- Sembrano JN, Polly DW Jr. How often is low back pain not coming from the back? Spine (Phila Pa 1976). 2009;34(1):E27-E32. doi:10.1097/ BRS.0b013e31818e0f5e.
- 8. Dreyfuss P, Dreyer SJ, Cole A, Mayo K. Sacroiliac joint pain. J Am Acad Orthop Surg. 2004;12(4):255–265.
- Laslett M, Young SB, Aprill CN, McDonald B. Diagnosing painful sacroiliac joints: a validity study of a McKenzie evaluation and sacroiliac provocation tests. Aust J Physiother. 2003;49(2):89–97.
- Vanaclocha V, Sáiz-Sapena N, Rivera-Paz M, Verdú-López F. Minimally invasive sacroiliac joint fusion: one-year and six-year follow-up. Neurosurgery. 2017;80(6):854-865. doi:10.1093/neuros/nyw113
- 11. Katz V, Schofferman J, Reynolds J. The sacroiliac joint: a potential cause of pain after lumbar fusion to the sacrum. Spine. 2003;28(10):S43-S48.
- 12. Maigne JY, Planchon CA. Sacroiliac joint pain after lumbar fusion. A study with anesthetic blocks. Eur Spine J. 2005;14(7):654-658. doi:10.1007/s00586-004-0692-6
- 13. DePalma MJ, Ketchum JM, Saullo TR. Etiology of chronic low back pain in patients having undergone lumbar fusion. Pain Med. 2011;12(5):732-739. doi:10.1111/j.1526-4637.2011.01098.x
- Liliang PC, Lu K, Liang CL, Tsai YD, Wang KW, Chen HJ. Sacroiliac joint pain after lumbar and lumbosacral fusion: findings using dual sacroiliac joint blocks. Pain Med. 2011;12(4):565-570. doi:10.1111/j.1526-4637.2011.01089.x
- Szadek KM, van der Wurff P, van Tulder MW, Zuurmond WW, Perez RS. Diagnostic validity of criteria for sacroiliac joint pain: a systematic review. J Pain. 2009;10(4):354-368. doi:10.1016/j.jpain.2008.09.014
- 16. Sturesson B, Selvik G, Udén A. Movements of the sacroiliac joints. A roentgen stereophotogrammetric analysis. Spine (Phila Pa 1976). 1989;14(2):162-165. doi:10.1097/00007632-198902000-00004

- 17. Jacob HA, Kissling RO. The mobility of the sacroiliac joints in healthy volunteers between 20 and 50 years of age. Clin Biomech (Bristol). 1995;10(7):352-361. doi:10.1016/0268-0033(95)00003-4
- 18. Wilke HJ, Fischer K, Jeanneret B, Claes L, Magerl F. In vivo measurement of 3-dimensionalmovementofth eiliosacraljoint[inGerman]. Z Orthop Ihre Grenzgeb. 1997;135(6):550-6.
- 19. Smidt GL, Wei SH, McQuade K, Barakatt E, Sun T. Sacroiliac motion for extreme hip positions: a fresh cadaver study. Spine (Phila Pa 1976). 1997;22(18):2073-2082. doi:10.1097/00007632-199709150-00007
- 20. Kiapour A, Joukar A, Elgafy H, Goel VK, Agarwal A, Benzel EC, et al. Relationship between limb length discrepancy and load distribution across the sacroiliac joint—A finite element study. J Orthop Res. 2012;30(10):1577–1580.
- 21. Fortin JD, Kissling RO, O'Connor BL, Vilensky JA. Sacroiliac joint innervation and pain. Am J Orthop (Belle Mead NJ). 1999;28(12):687–90.
- 22. Vilensky JA, O'Connor BL, Fortin JD, Merkel GJ, Xi M. Histologic analysis of the sacroiliac joint innervation in human subjects. Spine (Phila Pa 1976). 2002;27(11):1202–7.
- 23. Szadek KM, Hoogland PV, Zuurmond WW, de Lange JJ, Perez RS. Nociceptive nerve fibers in the sacroiliac joint in humans. Reg Anesth Pain Med. 2008Jan-Feb;33(1):36-43. PubMed PMID: 18155055.
- 24. Peretz AM, Hipp JA, Heggeness MH. The internal bony architecture of the sacrum. Spine (Phila Pa 1976). 1998;23(9):971-974. doi:10.1097/00007632-199805010-00001
- 25. Miller AN, Routt ML Jr. Variations in sacral morphology and implications for iliosacral screw fixation. J Am Acad Orthop Surg. 2012;20(1):8-16. doi:10.5435/ JAAOS-20-01-008
- 26. Hoel RJ, Ledonio CG, Takahashi T, Polly DW Jr. Sacral bone mineral density (BMD) assessment using opportunistic CT scans. J Orthop Res. 2017;35(1):160-166. doi:10.1002/jor.23362
- 27. Vleeming A, Schuenke MD, Masi AT, Carreiro JE, Danneels L, Willard FH. The sacroiliac joint: an overview of its anatomy, function and potential clinical implications. J Anat. 2012;221(6):537-567. doi:10.1111/j.1469-7580.2012.01564.x
- 28. Cohen, Steven P. MD. Sacroiliac Joint Pain: A Comprehensive Review of Anatomy, Diagnosis, and Treatment. Anesthesia & Analgesia 101(5):p 1440-1453, November 2005. | DOI: 10.1213/01. ANE.0000180831.60169.EA
- 29. Mabrouk OM, Ayad KE, Hady DAA. Contraction ratio of multifidus and erector spinae muscles in unilateral sacroiliac joint pain: a cross-sectional trial. Sci Rep. 2025;15:1730. doi:10.1038/s41598-024-84283-6
- Hermet M, Minichiello E, Flipo RM, Dubost JJ, Allanore Y, Ziza JM, Gaudin P, Thomas T, Dernis E, Glace B, Regnier A, Soubrier M. Infectious sacroiliitis: a retrospective, multicentre study of 39 adults. BMC Infect Dis. 2012 Nov 15;12:305. doi: 10.1186/1471-2334-12-305. PMID: 23153120; PMCID: PMC3519695.
- Slipman CW, Jackson HB, Lipetz JS, Chan KT, Lenrow DA, Vresilovic EJ. Sacroiliac joint pain referral zones. Arch Phys Med Rehabil. 2000;81(3):334–338.

- 32. Polly DW Jr. The Sacroiliac Joint: A Current State-ofthe-Art Review. JBJS Rev. 2024;12(2):e23.00151. Published 2024 Feb 5. doi:10.2106/JBJS.RVW.23.00151
- 33. Fortin JD, Dwyer AP, West SY, Pier J. Sacroiliac joint: pain referral maps upon applying a new injection/ arthrography technique. Spine (Phila Pa 1976). 1994;19(13):1475–82.
- 34. Fortin JD, Falco FJ. The Fortin finger test: an indicator of sacroiliac pain. Am J Orthop (Belle Mead NJ). 1997;26(7):477-480.
- Petersen T, Laslett M, Juhl C. Clinical classification in low back pain: best-evidence diagnostic rules based on systematic reviews. BMC Musculoskelet Disord. 2017;18(1):188. Published 2017 May 12. doi:10.1186/ s12891-017-1549-6
- Laslett M, Young SB, Aprill CN, McDonald B. Diagnosing painful sacroiliac joints: a validity study of a McKenzie evaluation and sacroiliac provocation tests. Aust J Physiother. 2003;49(2):89–97.
- Thawrani DP, Agabegi SS, Asghar F. Diagnosing Sacroiliac Joint Pain. J Am Acad Orthop Surg. 2019;27(3):85-93. doi:10.5435/JAAOS-D-17-00132
- Expert Panel on Neurological Imaging, Hutchins TA, Peckham M, et al. ACR Appropriateness Criteria[®] Low Back Pain: 2021 Update. J Am Coll Radiol. 2021;18(115):S361-S379. doi:10.1016/j.jacr.2021.08.002
- 39. Cohen AS, McNeill JM, Calkins E, Sharp JT, Schubart A. The "normal" sacroiliac joint. Analysis of 88 sacroiliac roentgenograms. Am J Roentgenol Radium Ther Nucl Med. 1967;100(3):559-563.
- 40. Eno JJ, Boone CR, Bellino MJ, Bishop JA. The prevalence of sacroiliac joint degeneration in asymptomatic adults. J Bone Joint Surg Am. 2015;97(11):932-936. doi:10.2106/JBJS.N.01101
- 41. Bhimreddy M, Weber-Levine C, Jiang K, et al. Sacroiliitis: current imaging modalities and future directions: a narrative review. Spine J. 2025;25(5):863-875. doi:10.1016/j.spinee.2024.11.011
- 42. Elgafy H, Semaan HB, Ebraheim NA, Coombs RJ. Computed tomography findings in patients with sacroiliac pain. Clin Orthop Relat Res. 2001;(382):112–118.
- 43. Mikula AL, Fogelson JL, Oushy S, Pinter ZW, Peters PA, Abode-Iyamah K, Sebastian AS, Freedman B, Currier BL, Polly DW, Elder BD. Change in pelvic incidence between the supine and standing positions in patients with bilateral sacroiliac joint vacuum signs. J NeurosurgSpine. 2021;34(4):617-22
- 44. Tsoi C, Griffith JF, Lee RKL, Wong PCH, Tam LS. Imaging of sacroiliitis: Current status, limitations and pitfalls. Quant Imaging Med Surg. 2019 Feb;9(2):318-335. doi: 10.21037/ qims.2018.11.10. PMID: 30976556; PMCID: PMC6414763.
- 45. Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. AJNR Am J Neuroradiol. 2015;36(4):811-816. doi:10.3174/ajnr.A4173
- Manchikanti L, Kaye AD, Soin A, et al. Comprehensive Evidence-Based Guidelines for Facet Joint Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines Facet Joint Interventions 2020 Guidelines. Pain Physician. 2020;23(3S):S1-S127.

- 47. Adams C, Banks KP. Bone Scan. In: StatPearls. Treasure Island (FL): StatPearls Publishing; August 14, 2023.
- 48. Rosenberg JM, Quint TJ, de Rosayro AM. Computerized tomographic localization of clinically-guided sacroiliac joint injections. Clin J Pain. 2000;16(1):18-21. doi:10.1097/00002508-200003000-00004
- 49. Broadhurst NA, Bond MJ. Pain provocation tests for the assessment of sacroiliac joint dysfunction. J Spinal Disord. 1998;11(4):341–345.
- 50. Dreyfuss P, Michaelsen M, Pauza K, McLarty J, Bogduk N. The value of medical history and physical examination in diagnosing sacroiliac joint pain. Spine (Phila Pa 1976). 1996;21(22):2594-2602. doi:10.1097/00007632-199611150-00009
- 51. Machado G C, Maher C G, Ferreira P H, Pinheiro M B, Lin C C, Day R O et al. Efficacy and safety of paracetamol for spinal pain and osteoarthritis: systematic review and meta-analysis of randomised placebo controlled trials BMJ 2015; 350 :h1225 doi:10.1136/ bmj.h1225
- 52. van der Gaag WH, Roelofs PD, Enthoven WT, van Tulder MW, Koes BW. Non-steroidal anti-inflammatory drugs for acute low back pain. Cochrane Database Syst Rev. 2020;4(4):CD013581. Published 2020 Apr 16. doi:10.1002/14651858.CD013581
- 53. Saragiotto BT, Machado GC, Ferreira ML, Pinheiro MB, Abdel Shaheed C, Maher CG. Paraceta- mol for low back pain. The Cochrane database of systematic reviews. 2016;2016(6):CD012230. doi:10.1002/14651858.CD012230
- 54. van Tulder MW, Touray T, Furlan AD, Solway S, Bouter LM. Muscle relaxants for non-specific low back pain. The Cochrane database of systematic reviews. 2003;2003(2):CD004252-CD004252. doi:10.1002/14651858.CD004252
- 55. Cashin AG, Folly T, Bagg MK, et al. Efficacy, acceptability, and safety of muscle relaxants for adults with non-specific low back pain: systematic review and meta-analysis. BMJ. 2021;374:n1446. doi:10.1136/bmj. n1446
- 56. Deyo RA, Von Korff M, Duhrkoop D. Opioids for low back pain. BMJ (Clinical research ed). 2015;350:g6380. doi:10.1136/bmj.g6380
- 57. Buynak R, Shapiro DY, Okamoto A, et al. Efficacy and safety of tapentadol extended release for the management of chronic low back pain: results of a prospective, randomized, double-blind, placebo- and active-controlled Phase III study. Expert opinion on pharmacotherapy. 2010;11(11):1787-1804. doi:10.1517 /14656566.2010.497720
- 58. Shaheed CA, Maher CG, Williams KA, Day R, McLachlan AJ. Efficacy, tolerability, and dose-dependent effects of opioid analgesics for low back pain a systematic review and meta-analysis. JAMA Internal Medicine. 2016;176(7):958-968. doi:10.1001/jamainternmed.2016.1251
- 59. Derry S, Wiffen PJ, Kalso EA, et al. Topical anal-gesics for acute and chronic pain in adults - an overview of Cochrane Reviews. The Cochrane database of systematic reviews. 2017;5(5):CD008609. doi:10.1002/14651858.CD008609.pub2

- 60. Voute M, Morel V, Pickering G. Topical Lidocaine for Chronic Pain Treatment. Drug design, development and therapy. 2021;15:4091-4103. doi:10.2147/DDDT. S328228
- 61. White WT, Patel N, Drass M, Nalamachu S. Lido- caine patch 5% with systemic analgesics such as gabapentin: a rational polypharmacy approach for the treatment of chronic pain. Pain medicine (Mal- den, Mass). 2003;4(4):321-330. doi:10.1111/j.1526-4637.2003.03045.x
- 62. Schmidt GL, Bhandutia AK, Altman DT. Management of sacroiliac joint pain. J Am Acad Orthop Surg. 2018;26(17):610-616. doi:10.5435/JAAOS-D-15-00063
- 63. Vleeming A, Buyruk HM, Stoeckart R, Karamursel S, Snijders CJ. An integrated therapy for peripartum pelvic instability: a study of the biomechanical effects of pelvic belts. Am J Obstet Gynecol. 1992;166(4):1243-1247. doi:10.1016/s0002-9378(11)90615-2
- 64. Sichting F, Rossol J, Soisson O, Klima S, Milani T, Hammer N. Pelvic belt effects on sacroiliac joint ligaments: a computational approach to understand therapeutic effects of pelvic belts. Pain Physician. 2014;17(1):43-51.
- 65. Roffey DM, Budiansky A, Coyle MJ, et al. Obesity and low back pain: is there a weight of evidence to support a positive relationship? Curr Obes Rep. 2013;2(3):241-250. doi:10.1007/s13679-013-0058-7
- 66. Chen LH, Weber K, Mehrabkhani S, et al. The effectiveness of weight loss programs for low back pain: a systematic review. BMC Musculoskelet Disord. 2022;23:488. doi:10.1186/s12891-022-05391-w
- Rashbaum, Ralph F. MD; Ohnmeiss, Donna D. Dr.Med.; Lindley, Emily M. PhD; Kitchel, Scott H. MD; Patel, Vikas V. MD. Sacroiliac Joint Pain and Its Treatment. Clinical Spine Surgery 29(2):p 42-48, March 2016. | DOI: 10.1097/BSD.00000000000359
- 68. Braun J, Bollow M, Seyrekbasan F, et al. Computed tomography guided corticosteroid injection of the sacroiliac joint in patients with spondyloarthropathy with sacroiliitis: clinical outcome and followup by dynamic magnetic resonance imaging. J Rheumatol. 1996;23(4):659-664.
- 69. Aydin SM, Gharibo CG, Mehnert M, Stitik TP. The role of radiofrequency ablation for sacroiliac joint pain: a meta-analysis. PM R. 2010;2(9):842-851. doi:10.1016/j. pmrj.2010.03.035
- Polly DW, Swofford J, Whang PG, et al. Randomized controlled trial of minimally invasive sacroiliac joint fusion using triangular titanium implants vs nonsurgical management for sacroiliac joint dysfunction: 12-month outcomes. Neurosurgery. 2015;77(5):674– 691. doi:10.1227/NEU.00000000000895
- Cher DJ, Frasco MA, Arnold RJ, Polly DW. Cost-effectiveness of minimally invasive sacroiliac joint fusion [published correction appears in Clinicoecon Outcomes Res. 2016 Jun 27;8:305. doi: 10.2147/CEOR. S107803.]. Clinicoecon Outcomes Res. 2015;8:1-14. Published 2015 Dec 18. doi:10.2147/CEOR.S94266

- 72. Duhon BS, Cher DJ, Wine KD, Lockstadt H, Kovalsky D, Soo CL. Safety and 6-month effectiveness of minimally invasive sacroiliac joint fusion: a prospective study. Med Devices (Auckl). 2013;6:219-229. Published 2013 Dec 13. doi:10.2147/MDER.S55197
- 73. Sturesson B, Udén A, Onsten I. Can an external frame fixation reduce the movements in the sacroiliac joint? A radiostereometric analysis of 10 patients. Acta Orthop Scand. 1999;70(1):42-46. doi:10.3109/17453679909000956
- 74. Duhon BS, Bitan F, Lockstadt H, et al. Triangular titanium implants for minimally invasive sacroiliac joint fusion: 2-year follow-up from a prospective multicenter trial. Int J Spine Surg. 2016;10:Article 13. doi:10.14444/3013
- 75. Rudolf L. Sacroiliac joint arthrodesis-MIS technique with titanium implants: report of the first 50 patients and outcomes. Open Orthop J. 2012;6:495–502. doi:10.2174/1874325001206010495
- 76. Whang PG, Duhon BS, Sasso RC, et al. Long-term prospective outcomes after minimally invasive sacroiliac joint fusion using triangular titanium implants. Med Devices (Auckl). 2019;12:411–422. doi:10.2147/MDER. S222351
- 77. Vanaclocha V, Herrera JM, Saiz-Sapena N, Rivera-Paz M, Verdu-Lopez F. Minimally invasive sacroiliac joint fusion, radiofrequency denervation, and conservative management for sacroiliac joint pain: 6-year comparative case series. Neurosurgery. 2018;82(1):48-55.
- 78. Martin CT, Haase L, Lender PA, Polly DW Jr. Minimally invasive sacroiliac joint fusion: the current evidence. Int J Spine Surg. 2020;14(Suppl 1):S20–S29. doi:10.14444/6072
- 79. Whang PG, Patel V, Duhon B, et al. Minimally Invasive SI Joint Fusion Procedures for Chronic SI Joint Pain: Systematic Review and Meta-Analysis of Safety and Efficacy. Int J Spine Surg. 2023;17(6):794-808. Published 2023 Dec 26. doi:10.14444/8543
- 80. Ivanov AA, Kiapour A, Ebraheim NA, Goel VK. Lumbar fusion leads to increases in angular motion and stress across sacroiliac joint: a finite element study. Spine (Phila Pa 1976). 2009;34(5):E162-E169. doi:10.1097/ BRS.0b013e3181978a35
- Polly DW Jr, Cher DJ, Wine KD, Whang PG, Frank CJ, Sembrano JN. Does level of response to SI joint block predict response to SI joint fusion? Int J Spine Surg. 2016;10:Article 4. doi:10.14444/3004
- 82. Eastlack RK, Soroceanu A, Mundis GM Jr, et al. Rates of Loosening, Failure, and Revision of Iliac Fixation in Adult Deformity Surgery. Spine (Phila Pa 1976). 2022;47(14):986-994. doi:10.1097/ BRS.00000000004356